

The universal ψ''/ψ' ratio as unambiguous signature for hard physics in nuclear reactions and in decays of beauty hadrons

L. Frankfurt^a, L. Gerland^a, M. Strikman^b, M. Zhalov^c

^a*School of Physics and Astronomy, Raymond and Beverly Sackler*

Faculty of Exact Science, Tel Aviv University, Ramat Aviv 69978, Tel Aviv , Israel

^b*Pennsylvania State University, University Park, Pennsylvania 16802*

^c*Petersburg Nuclear Physics Institute, Gatchina 188350, Russia*

February 1, 2008

Abstract

We suggest the ratio of the $\psi''(3770)/\psi'(3686)$ production cross sections as an unambiguous probe of the dominance of hard, point-like physics in charmonium production. If hard physics dominates the charmonium production the ratio : $\frac{\sigma(\psi'')}{\sigma(\psi')} \approx \frac{\Gamma(\psi'' \rightarrow l^+ l^-)}{\Gamma(\psi' \rightarrow l^+ l^-)} \approx 0.1$ is independent of the process. We argue that since the dominant $D\bar{D}$ decay mode of the ψ'' -meson can be detected with a high efficiency at the current facilities the effective counting rate for the ψ'' production should be comparable to that of the ψ' and hence it could be observed at the B -factories. Even in hadron induced processes the ratio will not be changed significantly by final state interactions due to the similar mass and size of these two charmonium states. A possible exception are the charmonium state interactions with low energy particles – comovers – which are especially relevant for the heavy ion collisions.

The production of the charmonium states in inclusive hadron-hadron collisions remains one of the challenges for the theory. Understanding of the charmonium production dynamics is important for a number of applications. This includes the dynamics of $B \rightarrow J/\psi + X$ decays which play an important

role in the investigations of the CP violation, see e.g. [3, 4]. The J/ψ production is also often considered as an effective probe of the heavy ion collisions. The matter is further complicated by the question whether the mass of the c -quark is sufficiently large as compared to the scale of the strong interaction to apply perturbation theory for the production of charm quark pairs in pp, pA, and AA collisions as well as in the decay of B -mesons.

So far experimental studies of charmonium states focused on the production of J/ψ , ψ' , and χ -states. The aim of this letter is to draw attention to the production of the ψ'' ($\psi(3770)$) meson as a method to probe the hardness of the process. Really in hard processes like in the process: $e\bar{e} \rightarrow$ charmonium state the contribution of S states should dominate the $c\bar{c}$ systems independently whether S or D states dominate in the wave function of charmonium state $\phi(k)$ in an average charmonium configuration. This is because the matrix element of the process is $\propto \int \phi(k) d^3k \propto \phi(r=0)$. But in a nonrelativistic charmonium model the wave function of a state with orbital momentum L e.g. the D wave is $\propto r^L$ at small relative interquark distances r because of the orbital momentum barrier. Thus the ratio of ψ' to ψ'' ($\psi(3770)$) should be the same for any process of hard production of charmonium states. The universality will be lost for the processes where the production will be influenced by nonperturbative QCD physics. So in this paper, we compare our results with the predictions of the nonrelativistic QCD (NRQCD) approach to the production of the onium states [1] which allowed to describe several features of the data. (Note that so far this approach meets problems in a number of cases - see ref. [2] for a recent review.) The NRQCD approach of ref. [17] assumed the dominance of D waves in the produced $c\bar{c}$ quark system which means nonhard production mechanisms play an important role. (The authors claims that the uncertainties in the evaluation of the relevant amplitudes may be one order of magnitude.) So we compare predictions of the NRQCD approach of ref. [17] and hard production mechanisms for the BELLE and the BABAR experiments.

We argue that the experimental observation of this process is feasible at the current facilities (which have a very good efficiency for detection of D -mesons), and would allow to check whether the ψ' and ψ'' are produced by a $c\bar{c}$ at short distances¹. Our suggestion is based on the well known

¹An important advantage of comparing ψ'' and ψ' production as compared to the case of J/ψ and ψ' is a near degeneracy of the masses of these states but different dominating decay channels.

observation that the properties of the ψ'' are well described (actually has been predicted) using the charmonium model by Eichten [5, 6]. Within this model it is found that the ψ'' is the $1D$ state with a small admixture of the $2S$ -wave and correspondingly ψ' is the $2S$ state with a small admixture of the $1D$ -wave. Namely

$$\begin{aligned} |\psi'\rangle &= \cos\theta |2S\rangle + \sin\theta |1D\rangle \quad , \\ |\psi''\rangle &= \cos\theta |1D\rangle - \sin\theta |2S\rangle \quad . \end{aligned} \quad (1)$$

Since only the S -wave contributes to the decay of ψ states into e^+e^- -pairs (at least in nonrelativistic charmonium models) the value of $\theta = 19 \pm 2^\circ$ can be determined from the data on the e^+e^- decay widths of ψ' and ψ'' as²

$$\tan^2(\theta) = \frac{\Gamma(\psi'' \rightarrow e^+e^-)}{\Gamma(\psi' \rightarrow e^+e^-)} \quad . \quad (2)$$

If the charmonium production is a short distance process, then it is proportional to the square of the wave function of charmonium at zero distance. Therefore, only the S -wave can contribute and the ψ''/ψ' ratio is universal, i.e. not energy dependent and not dependent on the process. Since the masses of ψ'' and ψ' are practically the same, the difference of the wave functions in the origin is the only source of the difference of the production rates. Correspondingly, the mixing model allows us to predict

$$\frac{\sigma(\psi'')}{\sigma(\psi')} = \tan^2(\theta) \approx 0.1 \quad . \quad (3)$$

It is instructive to consider the produced $c\bar{c}$ as a coherent superposition of the hadronic states. Since $m_{\psi'} \approx m_{\psi''}$ the coherence of the $c\bar{c}$ configuration which ultimately transforms into either ψ' or ψ'' holds up to the large distances $l_c \sim 2P/(m_{\psi''}^2 - m_{\psi'}^2) \approx \gamma \cdot 2.3 fm$, where γ is the γ -factor of the meson. l_c exceeds the size of the interaction region for all conceivable kinematics except deeply in the nucleus fragmentation region. Therefore it is more appropriate to think of the propagation of the $2S$ -state through the media with a subsequent decomposition of this state into ψ' and ψ'' .

²An estimate of ref. [6] within the charmonium model indicate that due to recoil corrections of the electromagnetic current the D-wave gives a $\sim 20\%$ contribution to the e.m. decay width. However, this model with the same parameters yields a too large electromagnetic decay width for the S-state.

The hadronic final state interactions cannot change significantly the predicted ratio. Indeed, soft QCD interactions cannot transform the S -state to D -state with significant probability. Significant values for the cross sections of the nondiagonal exclusive diffractive processes are a signature in the hadronic basis for the color screening phenomenon in the interaction of small color singlets [7] which should be a correction in soft QCD processes. The same conclusion is valid in the PQCD model for the charm dipole nucleon interactions. Hence the prediction of eq. (2) appears to be pretty robust provided ψ are produced via hard process³. At the same time if soft dynamics determines production of ψ'' , ψ' a stronger suppression of the ψ'' production compared to the ψ' production seems to be not natural since the charmonium models lead practically to the same sizes of the two mesons. For example, the model of ref. [5] gives for the average interquark distances $r_{2S} = 0.760$ fm and $r_{1D} = 0.786$ fm.

Let us consider the consequences of the prediction given by eq. (3) for the B -meson decays now. The branching fraction for B -decays into the $\psi(2S)$ was found by the BABAR-collaboration to be $0.297 \pm 0.020 \pm 0.020\%$ [3]. This branching fraction is averaged over the different B -mesons. The errors are statistical and systematical, respectively. The CLEO-collaboration found a comparable value of $0.316 \pm 0.014 \pm 0.023 \pm 0.016\%$ [4]. The third error is an overall scale factor. Due to the chosen decay channels of the $\psi(2S)$ ($\psi(2S) \rightarrow$ dileptons and $\psi(2S) \rightarrow J/\psi + \pi^+ + \pi^-$) we expect that these mesons are mostly ψ' states with a negligible admixture of ψ'' states. With the ratio of eq. (3) we predict the branching fractions for decays of the B -mesons into the ψ'' to be approximately 0.03%. In contrast the NRQCD approach predicts $B(B \rightarrow \delta_1 + X) = 0.28\%$ [17], where δ_1 is the D-wave charmonium with the quantum numbers of the ψ'' .

This branching fractions for decays of the B -mesons into the ψ'' seems to be smaller than the typical errors of these experiments. However, the statistics may not be problematic since in the measurement of the $B \rightarrow \psi(2S)$ both the BABAR and CLEO used the dileptonic decay channels of the $\psi(2S)$ or of the J/ψ which are smaller than 1% and 10%, respectively, while the decay of the ψ'' into D -mesons is the dominating channel. For this channel the detection efficiency is much higher. Hence it seems likely that

³Note that though the mass resolution in the current experiments is not good enough to resolve ψ' and ψ'' in the lepton mode the contribution of ψ'' into such decays is negligible due to much smaller value of $\Gamma_{e^+e^-}/\Gamma_{tot}$.

the counting rate for production of ψ'' via $D\bar{D}$ decays could be comparable to that for the ψ' decays via dilepton channels. If so this could provide a cross check of the measurement of the CP violation using new channels involving ψ'' production. In the refs. [8, 9] the collaborations CLEO and Belle showed that the channel $B^{\pm,0} \rightarrow \psi(2S) + K^{\pm,0}$ is 20% as large as the same channel for the J/ψ . However, due to the chosen decay channels this is the contribution of the ψ' only and the corresponding ψ'' may add another 20% if measured via the $D\bar{D}$ decays.

Additionally, the Belle Collaboration reported recently about the double charmonium production in e^+e^- annihilation at 10.6 GeV [10]⁴. In different theoretical works (one of the most recent is ref. [11]) is discussed if the cross section for double charmonium production can be described within a perturbative framework. However, the ratio of eq. (3) may be used here to test if these charmonium states result from local processes (e^+e^- annihilation, or PQCD) or from soft processes in the jet fragmentation. The BELLE collaboration found a production cross section for the ψ' of $\sigma(e^+e^- \rightarrow \psi' + X) = 0.67 \pm 0.09$ pb. The cross section of the ψ'' calculated in ref. [17] with the NRQCD approach is $\sigma(e^+e^- \rightarrow \delta_1 + X) = 0.043$ pb. This means that they predict roughly a factor of two more ψ'' than we, since this D-wave channel has the same order as the contribution of the S-wave.

In $p\bar{p}$ collisions at the Tevatron the CDF-collaboration found an inclusive production cross section for the ψ' of $\sigma(p\bar{p} \rightarrow \psi' + X) = 30.1 \pm 6.6^{+3.8}_{-4.2} \pm 6.6$ nb [12]. The errors are statistical, systematical, and due to the uncertainty of the $\psi' \rightarrow \mu^+\mu^-$ branching ratio⁵. Since the ψ' is measured here via the decay into $\mu^+\mu^-$ pairs, there is nearly no admixture of the ψ'' . Therefore we predict with eq. (3) an inclusive production cross section for the ψ'' of $\sigma(p\bar{p} \rightarrow \psi'' + X) \approx 3$ nb.

The ψ' meson production was also studied with 800 GeV protons at the Fermilab by the E789 and E772 collaboration using a set of nuclear targets [14, 15]. These data together with eq. (3) yield for the integrated over x_F and p_t production cross section of the ψ'' on a hydrogen target

$$\sigma(pp \rightarrow \psi'' + X) = 40 \pm 19 \text{ nb} \quad . \quad (4)$$

⁴We are grateful to A. Dumitru for drawing our attention to the refs. [9, 11] and for discussions.

⁵With the latest value for the $\psi' \rightarrow \mu^+\mu^-$ branching ratio of $\frac{\Gamma(\psi' \rightarrow \mu^+\mu^-)}{\Gamma(\psi' \rightarrow X)} = 0.007 \pm 0.0009$ [13] this would be $\sigma(p\bar{p} \rightarrow \psi' + X) = 33.1 \pm 6.6^{+3.8}_{-4.2} \pm 4.3$ nb.

Charmonium is also measured in proton nucleus collisions at a slightly higher energy ($E_{lab} = 920$ GeV) at HERA-B [16].

Note that several recent analyses of charmonium production processes performed within the NRQCD approach, see e.g. [17], have suggested the possibility of a significant production cross section of D-wave states. We compared our results for the BABAR and the BELLE experiment with the NRQCD predictions. According to NRQCD in these processes the direct contribution to the D-wave charmonium states is as large as the contribution to the ψ'' due to the admixture of the S-wave.

In this paper we compare the prediction of eq. (3) with the actual predictions of the NRQCD. From this angle it would be very important to look at the simplest process of the coherent photoproduction of charmonia. This is naturally a point-like process, in which the initial $c\bar{c}$ is produced in the S-state. Hence for this process all deviations from eq. (3) would be due to final state interactions. In particular it would be very interesting to study the A-dependence of the ψ''/ψ' ratio.

Also for heavy ion physics this could be an interesting investigations. The ratio of eq. (3) will not be changed due to hadronic final state interactions since these two states are very similar in size and mass. This can be tested at CERN-SPS by the NA60 collaboration [18] or at RHIC by the PHENIX collaboration [19]. Remember that for the inclusive charm production the current data both at FNAL energies [20] and at RHIC [21] are consistent with the linear dependence of the cross section on A which is expected within the QCD factorization theorem for the kinematics of the discussed experiments.

We discussed above that hadronic final state interactions will not change the ratio of eq. (3). However, in heavy ion collisions the formation of new phases is possible, e.g. the quark gluon plasma (QGP). It is not understood how to calculate this ratio in nonperturbative QCD in the QGP. However, for heavy ion collisions the thermal production of charmonium states is discussed in the literature [22]. In these models the ratio of the ψ'' and the ψ' yield is given by

$$\frac{\sigma(\psi'')}{\sigma(\psi')} = \left(\frac{m_{\psi''}}{m_{\psi'}} \right)^{3/2} \cdot \exp \left(\frac{m_{\psi''} - m_{\psi'}}{T} \right) \quad . \quad (5)$$

$T = 170 \div 180$ MeV in eq. (5) is the temperature, which is fitted to particle ratios in heavy ion collisions. Thus, eq. (5) yields in total

$$\frac{\sigma(\psi'')}{\sigma(\psi')} = 0.63 \div 0.65 \quad , \quad (6)$$

which is much larger than ratio of the cross sections predicted for the hard production processes. Moreover one should take into account that corrections to eq. (3) due to the higher ψ'' mass decrease this ratio even more. E.g. the pole mass of the propagators might yield a suppression factor of $\left(\frac{m_{\psi'}}{m_{\psi''}}\right)^2 = \left(\frac{3.686}{3.77}\right)^2 = 0.96$. The suppression which comes due to the shift of the initial partons to higher Bjorken x due to the higher mass yields for the gluon distribution a factor $\left(\frac{1-3.686/\sqrt{s}}{1-3.77/\sqrt{s}}\right)^{-12}$. For the relatively low SPS-energies of $\sqrt{s} = 20$ GeV this suppression factor is 0.94. However, since these corrections are smaller than 10% and since they even increase the difference between soft and hard production we can safely neglect these corrections here. Please note that for some experiments the kinematics can be chosen in a way that the mass difference between the ψ'' and the ψ' becomes unimportant, e.g. in hadron-hadron and hadron-nucleus collisions. Also, the mass difference becomes unimportant at high transverse momenta, since here the transverse mass not the rest mass of the particles is the relevant parameter.

To summarize, we have presented arguments in favour of a significant rate of the ψ'' production in various hard processes and for the sensitivity of the ratio of the ψ''/ψ' rate to the dynamics of charmonium production. A number of the current experiments can answer this question by scanning their existing data.

Acknowledgement:

We thank E. Braaten and T. Ullrich for useful discussions. This work was supported in part by GIF and DOE. L.G. thanks the Minerva Foundation for support.

References

- [1] G. T. Bodwin, E. Braaten and G. P. Lepage, Phys. Rev. D **51**, 1125 (1995) [Erratum-ibid. D **55**, 5853 (1997)] [arXiv:hep-ph/9407339].
- [2] G. T. Bodwin, arXiv:hep-ph/0212203.
- [3] B. Aubert *et al.* [BABAR Collaboration], arXiv:hep-ex/0207097.
- [4] S. Anderson *et al.* [CLEO Collaboration], arXiv:hep-ex/0207059.

- [5] E. Eichten, K. Gottfried, T. Kinoshita, J. B. Kogut, K. D. Lane and T. M. Yan, Phys. Rev. Lett. **34** (1975) 369 [Erratum-ibid. **36** (1976) 1276].
- [6] J. M. Richard, Z. Phys. C **4** (1980) 211.
- [7] L. Frankfurt and M. Strikman, Prog. Part. Nucl. Phys. **27**, 135 (1991).
- [8] G. Bonvicini *et al.* [CLEO Collaboration], Phys. Rev. Lett. **84**, 5940 (2000) [arXiv:hep-ex/0003004].
- [9] K. Abe *et al.* [BELLE Collaboration], arXiv:hep-ex/0211047.
- [10] K. Abe *et al.* [Belle Collaboration], Phys. Rev. Lett. **89**, 142001 (2002) [arXiv:hep-ex/0205104].
- [11] G. T. Bodwin, J. i. Lee and E. Braaten, arXiv:hep-ph/0212352.
- [12] F. Abe *et al.* [CDF Collaboration], Phys. Rev. Lett. **69**, 3704 (1992).
- [13] K. Hagiwara *et al.* [Particle Data Group Collaboration], Phys. Rev. D **66**, 010001 (2002).
- [14] M. H. Schub *et al.* [E789 Collaboration], Phys. Rev. D **52**, 1307 (1995) [Erratum-ibid. D **53**, 570 (1996)].
- [15] D. M. Alde *et al.*, Phys. Rev. Lett. **66**, 133 (1991).
- [16] I. Abt *et al.* [HERA-B Collaboration], arXiv:hep-ex/0211033.
- [17] L. K. Hao, K. Y. Liu and K. T. Chao, Phys. Lett. B **546**, 216 (2002) [arXiv:hep-ph/0206226].
F. Yuan, C. F. Qiao and K. T. Chao, Phys. Rev. D **56**, 329 (1997) [arXiv:hep-ph/9701250].
P. w. Ko, J. Lee and H. S. Song, Phys. Lett. B **395**, 107 (1997) [arXiv:hep-ph/9701235].
- [18] B. Lenkeit [NA60-Collaboration], arXiv:nucl-ex/0108015.
- [19] K. Adcox *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **88**, 192303 (2002) [arXiv:nucl-ex/0202002].

- [20] P. L. McGaughey, J. M. Moss and J. C. Peng, *Ann. Rev. Nucl. Part. Sci.* **49** (1999) 217 [arXiv:hep-ph/9905409].
- [21] D. G. D'Enterria [PHENIX Collaboration], arXiv:hep-ex/0205060.
- [22] M. I. Gorenstein, A. P. Kostyuk, L. McLerran, H. Stöcker and W. Greiner, *J. Phys. G* **28**, 2151 (2002).
P. Braun-Munzinger and J. Stachel, *Phys. Lett. B* **490**, 196 (2000) [arXiv:nucl-th/0007059].
R. L. Thews, M. Schroedter and J. Rafelski, *Phys. Rev. C* **63**, 054905 (2001) [arXiv:hep-ph/0007323].